arXiv:1303.2485v1 [math.OA] 11 Mar 2013

STRONGLY IRREDUCIBLE OPERATORS AND INDECOMPOSABLE REPRESENTATIONS OF QUIVERS ON INFINITE-DIMENSIONAL HILBERT SPACES

MASATOSHI ENOMOTO AND YASUO WATATANI

Abstract. We study several classes of indecomposable representations of quivers on infinite-dimensional Hilbert spaces and their relation. Many examples are constructed using strongly irreducible operators. Some problems in operator theory are rephrased in terms of representations of quivers. We shall show two kinds of constructions of quite non-trivial indecomposable Hilbert representations (H, f) of the Kronecker quiver such that $End(H, f) = \mathbb{C}I$ which is called transitive. One is a perturbation of a weighted shift operator by a rank-one operator. The other one is a modification of an unbounded operator used by Harrison,Radjavi and Rosenthal to provide a transitive lattice.

KEYWORDS: strongly irreducible operators, quiver, indecomposable representation, Hilbert space.

AMS SUBJECT CLASSIFICATION: Primary 47A65, Secondary 46C07,47A15, 15A21, 16G20, 16G60.

1. Introduction

A bounded operator T on a Hilbert space H is said to be strongly irreducible if T cannot be decomposed to a non-trivial (not necessarily orthogonal) direct sum of two operators, that is, if there exist no non-trivial invariant (closed) subspaces M and N of T such that $M \cap N = 0$ and $M + N = H$. A strongly irreducible operator is an infinite-dimensional analog of a Jordan block. The notion of strongly irreducible operator was introduced by F. Gilfeather in [Gi]. We refer to good monographs [JW1] and [JW2] by Jiang and Wang on strongly irreducible operators.

On the other hand Gabriel [Ga] introduced a finite-dimensional (linear) representations of quivers by attaching vector spaces and linear maps for vertices and edges of quivers respectively. A finite-dimensional indecomposable representation of a quiver is a direct graph generalization of a Jordan block. We regard indecomposable representation of a quiver on a Hilbert space as an infinite-dimensional generalization of both a Jordan block and a finite-dimensional indecomposable representation of a quiver. We study several classes of indecomposable representations of quivers on infinite-dimensional Hilbert spaces and their relation. Many examples of indecomposable representations of quivers are constructed using strongly irreducible operators. Moreover some problems in operator theory are rephrased in terms of representations of quivers.

Remember that we studied the relative positions of subspaces in a separable infinite-dimensional Hilbert space in [EW1] after Nazarova [Na1], Gelfand and Ponomarev [GP]. We shall describe a close relation between the relative positions of subspaces in Hilbert spaces and Hilbert representations of quivers in the final section.

In our paper we only need the beginning of the theory of representations of quivers on finite-dimensional vector space, for example, see Bernstein-Gelfand-Ponomarev [BGP], Donovan-Freislish [DF], V. Dlab-Ringel [DR], Gabriel-Roiter [GR], Kac [Ka], Nazarova [Na2] ....

We should remark that locally scalar representations of quivers in the category of Hilbert spaces were introduced by Kruglyak and Roiter [KRo]. They associate operators and their adjoint operators with arrows and classify them up to the unitary equivalence. They proved an analog of Gabriel's theorem. Their study is connected with representations of *-algebras generated by linearly related orthogonal projections , see for example, S. Kruglyak, V. Rabanovich and Y. Samoilenko [KRS] and Y. P. Moskaleva and Y. S. Samoilenko [MS].

In [EW3], we constructed an indecomposable infinite-dimensional Hilbert representation for any quiver whose underlying undirected graph is one of extended Dynkin diagrams $\tilde{A_n}$ $(n \geq 0)$, $\tilde{D_n}$ $(n \geq 4)$, $\tilde{E_6}$,$\tilde{E_7}$ and $\tilde{E_8}$, using the unilateral shift S. If we replace the unilateral shift S there by any strongly irreducible operator, then the corresponding Hilbert representation is still indecomposable by the same calculation. This fact also suggests us to use strong irreducible operators to construct indecomposable Hilbert representations of quivers.

We recall infinite-dimensional representations in purely algebraic setting. In [Au] Auslander found that if a finite- dimensional algebra is not of finite representation type, then there exist indecomposable modules which are not of finite length. Ringel [Ri1] developed a general theory of infinite-dimensional representations of tame, hereditary algebra. There exist many works after them and they form an active area of research in representation theory of algebras.

In our paper we study infinite-dimensional Hilbert (space) representations of quivers using operator theory. We note that there exist subtle difference among purely algebraic infinite-dimensional representations of quivers, infinite-dimensional Banach (space) representations of quivers and infinite-dimensional Hilbert (space) representations of quivers. We also note that the analytic aspect of Hardy space is quite important in our setting. For example, if the endomorphism algebra of a Hilbert representation of a quiver is isomorphic to the Hardy algebra $H^\infty(\mathbb{T})$, then the representation is indecomposable, because the the Hardy algebra $H^{\infty}(\mathbb{T})$ has no non-trivial idempotents by the F. and M. Riesz Theorem. This is indeed the case of the Hilbert representation corresponding to the unilateral shift operator. In this way we believe that the analytic operator algebra theory will come in here.

We shall show two kinds of constructions of quite non-trivial indecomposable Hilbert representation (H, f) of the Kronecker quiver such that $End(H, f) = \mathbb{C}I$ which is called transitive. One is a perturbation of a weighted shift operator by a rank-one operator. This is an analogue of a construction of indecomposable representations using linear functionals on the space $K(x)$ of rational functions over an algebraically closed field K studied in representation theory of algebras, for example, see Ringel [Ri2], Fixmann [Fi], Okoh [Ok] and Dean- Zorzitto [DZ]. We replace the rational function field $K(x)$ by Hardy spaces $H^{\infty}(\mathbb{T})$ or $H^2(\mathbb{T})$ properly in our setting . We have an analogy of ring extension between $(\mathbb{C}[x] \subset \mathbb{C}(x))$ and $(\mathbb{C}[x] \subset H^{\infty}(\mathbb{T}))$ in mind. Our analogy is supported by an important fact that both the rational function field $\mathbb{C}(x)$ and the Hardy algebra $H^{\infty}(\mathbb{T})$ have no non-trivial idempotents. But we warn the readers of subtle differences among them.

The other construction of transitive representations of the Kronecker quiver is given by a modification of an unbounded operator used by Harrison,Radjavi and Rosenthal [HRR] to provide a transitive lattice.

2. Hilbert representations of quivers

A quiver $\Gamma = (V, E, s, r)$ is a quadruple consisting of the set V of vertices, the set E of arrows, and two maps $s, r : E \to V$, which associate with each arrow $\alpha \in E$ its support $s(\alpha)$ and range $r(\alpha)$. We sometimes denote by $\alpha : x \to y$ an arrow with $x = s(\alpha)$ and $y = r(\alpha)$. Thus a quiver is just a directed graph. We denote by $|\Gamma|$ the underlying undirected graph of a quiver Γ. A quiver Γ is said to be connected if $|\Gamma|$ is a connected graph. A quiver Γ is said to be finite if both V and E are finite sets. A path of length m is a finite sequence $\alpha = (\alpha_1, \dots, \alpha_m)$ of arrows such that $r(\alpha_k) = s(\alpha_{k+1})$ for $k = 1, \dots, m-1$. Its support is $s(\alpha) = s(\alpha_1)$ and its range is $r(\alpha) = r(\alpha_m)$. A path of length $m \geq 1$ is called a *cycle* if its support and range coincide. A cycle of length one is called a *loop*. A quiver is called *acyclic* if it contains no cycles.

Definition. Let $\Gamma = (V, E, s, r)$ be a finite quiver. We say that (H, f) is a *Hilbert representation* of Γ if $H = (H_v)_{v \in V}$ is a family of Hilbert spaces and $f = (f_\alpha)_{\alpha \in E}$ is a family of bounded linear operators such that $f_\alpha : H_{s(\alpha)} \to H_{r(\alpha)}$ for $\alpha \in E$.

Definition. Let $\Gamma = (V, E, s, r)$ be a finite quiver. Let (H, f) and (K, g) be Hilbert representations of Γ. A *homomorphism* $T : (H, f) \to$

(K, g) is a family $T = (T_v)_{v \in V}$ of bounded operators $T_v : H_v \to K_v$ satisfying, for any arrow $\alpha \in E$

$$T_{r(\alpha)} f_\alpha = g_\alpha T_{s(\alpha)}.$$

The composition $T \circ S$ of homomorphisms T and S is defined by the composition of operators: $(T \circ S)_v = T_v S_v$ for $v \in V$. Thus we have obtained a category $HRep(\Gamma)$ of Hilbert representations of Γ

We denote by $Hom((H, f), (K, g))$ the set of homomorphisms $T : (H, f) \to (K, g)$. We denote by $End(H, f) := Hom((H, f), (H, f))$ the set of endomorphisms. Then we can regard $End(H, f)$ as a subalgebra of $\oplus_{v \in V} B(H_v)$.

In the paper we carefully distinguish the following two classes of operators. A bounded operator A on a Hilbert space is called an *idempotent* if $A^2 = A$ and A is said to be a *projection* if $A^2 = A = A^*$. We denote by

$$\begin{aligned} Idem(H, f) &:= \{T \in End(H, f) \mid T^2 = T\} \\ &= \{T = (T_v)_{v \in V} \in End(H, f) \mid T_v^2 = T_v \ (\text{for any } v \in V)\} \end{aligned}$$

the set of idempotents of $End(H, f)$. Let $0 = (0_v)_{v \in V}$ be a family of zero operators 0_v and $I = (I_v)_{v \in V}$ be a family of identity operators I_v. The both endomorphisms 0 and I are in $Idem(H, f)$.

Let $\Gamma = (V, E, s, r)$ be a finite quiver and (H, f), (K, g) be Hilbert representations of Γ. We say that (H, f) and (K, g) are *isomorphic*, denoted by $(H, f) \cong (K, g)$, if there exists an isomorphism $\varphi : (H, f) \to (K, g)$, that is, there exists a family $\varphi = (\varphi_v)_{v \in V}$ of bounded invertible operators $\varphi_v \in B(H_v, K_v)$ such that, for any arrow $\alpha \in E$,

$$\varphi_{r(\alpha)} f_\alpha = g_\alpha \varphi_{s(\alpha)}.$$

Hilbert representations (H, f) and (K, g) of Γ are said to be *relatively prime* if $Hom((H, f), (K, g)) = 0$ and $Hom((K, g), (H, f)) = 0$. If two non-zero Hilbert representations (H, f) and (K, g) are relatively prime, then they are not isomorphic.

We say that (H, f) is a finite-dimensional representation if H_v is finite-dimensional for all $v \in V$. And (H, f) is an infinite-dimensional representation if H_v is infinite-dimensional for some $v \in V$.

We shall recall a notion of indecomposable representation in [EW3], that is, a representation which cannot be decomposed into a direct sum of smaller representations anymore.

Definition.(Direct sum) Let $\Gamma = (V, E, s, r)$ be a finite quiver. Let (K, g) and (K', g') be Hilbert representations of Γ. Define the direct sum $(H, f) = (K, g) \oplus (K', g')$ by

$$H_v = K_v \oplus K'_v \ (\text{ for } v \in V) \quad \text{and} \quad f_\alpha = g_\alpha \oplus g'_\alpha \ (\text{ for } \alpha \in E).$$

We say that a Hilbert representation (H, f) is zero, denoted by $(H, f) = 0$, if $H_v = 0$ for any $v \in V$.

Definition.(Indecomposable representation) A Hilbert representation (H, f) of Γ is called *decomposable* if (H, f) is isomorphic to a direct sum of two non-zero Hilbert representations. A non-zero Hilbert representation (H, f) of Γ is said to be *indecomposable* if it is not decomposable, that is, if $(H, f) \cong (K, g) \oplus (K', g')$ then $(K, g) \cong 0$ or $(K', g') \cong 0$.

The following proposition is used frequently to show the indecomposability in concrete examples.

Proposition 2.1. ([EW3, Proposition 3.1.]) Let (H, f)be a Hilbert representation of a quiver Γ. Then the following conditions are equivalent:

(1) (H, f) is indecomposable.
(2) $Idem(H, f) = \{0, I\}$.

Remark. (H, f) is decomposable if and only if there exist families $K = (K_x)_{x \in V}$ and $K' = (K'_x)_{x \in V}$ of closed subspaces K_x and K'_x of H_x with $K_x \cap K'_x = 0$ and $K_x + K'_x = H_x$ satisfying K and K' are non-zero such that $f_\alpha K_x \subset K_y$ and $f_\alpha K'_x \subset K'_y$ for any arrow $\alpha : x \to y$.

The indecomposability of Hilbert representations of a quiver is an isomorphic invariant, but it is *not* a unitarily equivalent invariant. Therefore we cannot replace the set $Idem(H, f)$ of idempotents of endomorphisms by the subset of idempotents of endomorphisms which are consists of projections to show the indecomposability.

Example 1. Let $H_0 = \mathbb{C}^2$. Fix an angle θ with $0 < \theta < \pi/2$. Consider one-dimensional subspaces $H_1 = \mathbb{C}(1, 0)$ and $H_2 = \mathbb{C}(\cos\theta, \sin\theta)$ of H_0 spanned by vectors $(1, 0)$ and $(\cos\theta, \sin\theta)$ in H_0. Consider the following quiver Γ :

$$\begin{tikzcd} \circ_1 \arrow[r, "\alpha_1"] & \circ_0 & \circ_2 \arrow[l, "\alpha_2"'] \end{tikzcd}$$

Define a Hilbert representation (H, f) of Γ by $H = (H_i)_{i=0,1,2}$ and canonical inclusion maps $f_i = f_{\alpha_i} : H_i \to H_0$ for $i = 1, 2$. Then the Hilbert representation (H, f) is decomposable. But if an idempotent $P = (P_v)_{v \in V} \in End(H, f)$ satisfies that P_v is a projection for any $v \in V$, then $P = 0$ or $P = I$. In fact, since $H_0 = H_1 + H_2$ and $H_1 \cap H_2 = 0$, for any $x \in H_0$, there exist unique $x_1 \in H_1$ and $x_2 \in H_2$ such that $x = x_1 + x_2$. There exists an idempotent $T_0 : H_0 \to H_0$ such that $T_0 x = x_1$. Put $T_1 = id : H_1 \to H_1$ and $T_2 = 0 : H_2 \to H_2$. Then $T = (T_i)_{i=0,1,2}$ is an idempotent in $End(H, f)$ such that $T \neq 0$ and $T \neq I$. Hence (H, f) is decomposable. But take any idempotent $P = (P_v)_{v \in V} \in End(H, f)$ such that P_v is a projection for any $v \in V$. Then $P_0 H_1 \subset H_1$ and $P_0 H_2 \subset H_2$. Let E_i be the projection of H_0 onto H_i for $i = 1, 2$. Since P_0 is self adjoint, P_0 commutes with E_1 and E_2. Since the C^*-algebra generated by E_1 and E_2 is $B(H_0) = M_2(\mathbb{C})$ and P_0 is a projection, $P_0 = 0$ or $P_0 = I$. If $P_0 = 0$, then $P_1 = 0$ and $P_2 = 0$. Similarly if $P_0 = I$, then $P_1 = I$ and $P_2 = I$. Hence $P = 0$

or $P = I$. We remark that the system $(H_0; H_1, H_2)$ of two subspaces is isomorphic to

$$(\mathbb{C}^2; \mathbb{C} \oplus 0, 0 \oplus \mathbb{C}) \cong (\mathbb{C}; \mathbb{C}, 0) \oplus (\mathbb{C}; 0, \mathbb{C}).$$

Hence the relative position of two subspaces $(H_0; H_1, H_2)$ is decomposable. See [EW3, Remark after Proposition 3.1.].

Definition. A non-zero Hilbert representation (H, f) of a quiver Γ is called *irreducible* if $P = (P_v)_{v \in V} \in End(H, f)$ is an idempotent and P_v is a projection for any $v \in V$, then $P = 0$ or $P = I$. A non-zero Hilbert representation (H, f) is not irreducible if and only if there exist families $K = (K_x)_{x \in V}$ and $K' = (K'_x)_{x \in V}$ of closed subspaces K_x and K'_x of H_x with $K_x \perp K'_x$ and $K_x + K'_x = H_x$ satisfying K and K' are non-zero such that $f_\alpha K_x \subset K_y$ and $f_\alpha K'_x \subset K'_y$ for any arrow $\alpha : x \to y$. For example, the Hilbert representation (H, f) in Example 1 above is irreducible but is not indecomposable. We should be careful that irreducibility is a unitarily invariant notion and not a isomorphically invariant notion.

We recall the following elementary but fundamental relation between Hilbert representation theory of quivers and single operator theory:

Theorem 2.2 ([EW3]). *Let L_1 be one-loop quiver, so that L_1 has one vertex 1 and one arrow $\alpha : 1 \to 1$. The underlying undirected graph is an extended Dynkin diagram $\tilde{A}_0$. For a bounded operator A on a Hilbert space H, consider a Hilbert representation (H^A, f^A) of L_1 such that $H_1^A = H$ and $f_\alpha^A = A$. Then the Hilbert representation (H^A, f^A) is indecomposable if and only if A is strongly irreducible. The endomorphism ring $End(H, f)$ can be identified with the commutant $\{A\}'$. Moreover two bounded operators A and B are similar if and only if the corresponding Hilbert representations (H^A, f^A) and $(H^B, f^B)$are isomorphic.*

Therefore it is fruitful to regard the study of indecomposable Hilbert representations of general quivers as a generalization of the study of strongly irreducible operators.

Example 2. Consider one-loop quiver L_1 as above. Let $H = \ell^2(\mathbb{N})$ and $S \in B(H)$ the unilateral shift. Then (H^S, f^S) is indecomposable.

Definition. Let $\Gamma = (V, E, s, r)$ be a finite quiver and (H, f) a Hilbert representation of Γ. A Hilbert representation (K, g) of Γ is called a *subrepresentation* of (H, f) if for any vertex $v \in V$, K_v is a (closed) subspace of H_v and for any edge $\alpha \in E$, $g_\alpha = f_\alpha|_{K_{s(\alpha)}}$. In particular we have $f_\alpha(K_{s(\alpha)}) \subset K_{r(\alpha)}$.

Definition. A non-zero Hilbert representation (H, f) of a quiver Γ is called *simple* if (H, f) has only trivial subrepresentations 0 and (H, f).

A Hilbert representation (H, f) of Γ is called *canonically simple* if there exists a vertex $v_0 \in V$ such that $H_{v_0} = \mathbb{C}$, $H_v = 0$ for any other vertex $v \neq v_0$ and $f_\alpha = 0$ for any $\alpha \in E$. It is clear that if a Hilbert representation (H, f) of Γ is canonically simple, then (H, f) is simple. It is trivial that if a Hilbert representation (H, f) of Γ is simple, then (H, f) is indecomposable.

We can rephrase the invariant subspace problem in terms of simple representations of a one-loop quiver. Let L_1 be one-loop quiver, so that L_1 has one vertex 1 and one arrow $\alpha : 1 \to 1$. Any bounded operator A on a non-zero Hilbert space H gives a Hilbert representation (H^A, f^A) of L_1 such that $H_1^A = H$ and $f_\alpha^A = A$. Then the operator A has only trivial invariant subspaces if and only if the Hilbert representation (H^A, f^A) of L_1 is simple. If H is one-dimensional and A is a non-zero scalar operator, then the Hilbert representation (H^A, f^A) of L_1 is simple but is not canonically simple. If H is finite-dimensional with $\dim H \geq 2$, then the Hilbert representation (H^A, f^A) of L_1 is not simple, because any operator A on H has a non-trivial invariant subspace. If H is countably infinite-dimensional, then we do not know whether the Hilbert representation (H^A, f^A) of L_1 is not simple. In fact this is the invariant subspace problem, that is, the question whether any operator A on H has a non-trivial (closed) invariant subspace.

Definition. A Hilbert representation (H, f) of a quiver Γ is called *transitive* if $End(H, f) = \mathbb{C}I$. It is clear that if a Hilbert representation (H, f) is canonically simple, then (H, f) is transitive. If a Hilbert representation (H, f) of Γ is transitive , then (H, f) is indecomposable. In fact, since $End(H, f) = \mathbb{C}I$, any idempotent endomorphism T is 0 or I. In purely algebraic setting, a representation of a quiver is called a *brick* if its endomorphism ring is a division ring. But for a Hilbert representation (H, f) of a quiver, $End(H, f)$ is a Banach algebra and not isomorphic to its purely algebraic endomorphism ring in general, because we only consider bounded endomorphisms. By Gelfand-Mazur theorem, any Banach algebra over $\mathbb{C}$ which is a division ring must be isomorphic to $\mathbb{C}$. Therefore the reader may use "brick" instead of "transitive Hilbert representation" if he does not confuse the difference between purely algebraic endomorphism ring and $End(H, f)$.

Remark. A lattice $\mathcal{L}$ of subspaces of a Hilbert space H containing 0 and H is called a transitive lattice if

$$\{A \in B(H) \mid AM \subset M \text{ for any } M \in \mathcal{L}\} = \mathbb{C}I.$$

See, for example, Radjavi-Rosenthal [RR, 4.7.].

Let $\mathcal{L} = \{0, M_1, M_2, \ldots, M_n, H\}$ be a finite lattice. Consider a n subspace quiver $R_n = (V, E, s, r)$, that is, $V = \{1, 2, \ldots, n, n+1\}$ and $E = \{\alpha_k \mid k = 1, \ldots, n\}$ with $s(\alpha_k) = k$ and $r(\alpha_k) = n+1$ for $k = 1, \ldots, n$. Then there exists a Hilbert representation (K, f) of R_n

such that $K_k = M_k$, $K_{n+1} = H$ and $f_{\alpha_k} : M_k \to H$ is an inclusion for $k = 1, \dots, n$. Then the lattice $\mathcal{L}$ is transitive if and only if the corresponding Hilbert representation (H, f) is transitive. This fact guarantees the terminology "transitive" in the above.

Example 3. Let L_2 be 2-loop, that is, L_2 is a quiver with one vertex $\{v\}$ and two loops $\{\alpha, \beta\}$. Let S be the unilateral shift on $\ell^2(\mathbb{N})$. Define a Hilbert representation (H, f) of L_2 by $H_v = \ell^2(\mathbb{N})$ and $f_\alpha = S, f_\beta = S^*$. Then (H, f) is simple and transitive but (H, f) is not canonically simple. In fact, since $End(H, f)$ is given by the commutant $\{S, S^*\}'$ and $\{S, S^*\}' = \mathbb{C}I$, (H, f) is transitive. Any subrepresentation of (H, f) is given by the common invariant subspaces of S and S^*, which is 0 or $\ell^2(\mathbb{N})$. Hence any subrepresentation of (H, f) is 0 or (H, f).



Example 4. Let $K_3 = (V, E, s, r)$ be a 3-Kronecker quiver, so that $V = \{1, 2\}$, $E = \{\alpha_1, \alpha_2, \alpha_3\}$ and $s(\alpha_i) = 1, r(\alpha_i) = 2$ for $i = 1, 2, 3$. Let S be the unilateral shift on $\ell^2(\mathbb{N})$. Define a Hilbert representation (H, f) by $H_1 = H_2 = \ell^2(\mathbb{N})$ and $f_1 = S$, $f_2 = S^*$, $f_3 = I$. Then the Hilbert representation (H, f) is simple but not canonically simple.

Example 5. A bounded operator$A$ on a Hilbert space H is called *unicellular* if the lattice of invariant subspaces of A is totally ordered. See [RR] for unicellular operators. For example, the unilateral shift S is unicellular. Any non-zero unicellular operator is strongly irreducible. Let L_1 be one-loop quiver, so that L_1 has one vertex 1 and one arrow $\alpha : 1 \to 1$. Consider a Hilbert representation (H^A, f^A) of L_1 such that $H^A_1 = H$ and $f^A_\alpha = A$. If A is unicellular, then Hilbert representation (H^A, f^A) of L_1 is indecomposable. Let L_2 be 2-loop, that is, L_2 is a quiver with one vertex $\{v\}$ and two loops $\{\alpha, \beta\}$. Define a Hilbert representation (H, f) of L_2 by $H_v = H$ and $f_\alpha = A, f_\beta = A^*$. If A is a unicellular operator, then (H, f) is simple. In fact any subrepresentation (K, g) is given by a common invariant subspace M of A and A^*. Then M and $M^\perp$ is an invariant subspace of A. Since A is unicellular, $M = 0$ or $M = H$.

We summarize relations among several classes of Hilbert representations of quivers.



Example 6. Consider 2-loop quiver L_2, that is, L_2 is a quiver with one vertex $\{v\}$ and two loops $\{\alpha, \beta\}$. Define $H_v = \mathbb{C}^2$ and

$$f_\alpha = A := \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad f_\beta = B := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Then the Hilbert representation (H, f) is transitive but is not simple. Since

$$End(H, f) = \{T \in M_2(\mathbb{C}) \mid TA = AT \text{ and } TB = BT\} = \mathbb{C}I,$$

(H, f) is transitive. Define $K_v = \mathbb{C} \oplus 0$ and $g_\alpha = id_{K_v}$ and $g_\beta = 0$. Then (K, g) is a subrepresentation such that $(K, g) \neq 0$ and $(K, g) \neq (H, f)$. Therefore (H, f) is not simple.

Example 7. Consider 2-loop quiver L_2, that is, L_2 is a quiver with one vertex $\{v\}$ and two loops $\{\alpha, \beta\}$. Define $H_v = \mathbb{C}^2$ and

$$f_\alpha = A := \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad f_\beta = C := \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Then the Hilbert representation (H, f) is transitive and simple. In fact since

$$End(H, f) = \{T \in M_2(\mathbb{C}) \mid TA = AT \text{ and } TC = CT\} = \mathbb{C}I,$$

(H, f) is transitive. Let (K, g) be subrepresentation of (H, f). Since K_v is a common invariant subspace for A and C, K_v is 0 or $H_v = \mathbb{C}^2$. Hence $(K, g) = 0$ or $(K, g) = (H, f)$. Thus (H, v) is simple.

We collect some elementary facts:

Proposition 2.3. *Let (H, f) be a finite-dimensional Hilbert representation of a quiver Γ. If (H, f) is simple, then (H, f) is transitive.*

Proof. Assume that (H, f) is not transitive. Then the following two cases occur:
(A) There exists a vertex $u \in V$ and an endomorphism $T \in End(H, f)$ such that $T_u \notin \mathbb{C}I_u$.
(B) There exist vertices $v_1 \neq v_2$ and scalars $\lambda_1 \neq \lambda_2$ such that $T_{v_1} = \lambda_1 I_{v_1}$, $T_{v_2} = \lambda_2 I_{v_2}$ with $H_{v_1} \neq 0$ and $H_{v_2} \neq 0$.
In either case we shall show that there exists a non-trivial subrepresentation.
The case (A): There exists a scalar λ such that the eigenspace $0 \neq \mathrm{Ker}(T_u - \lambda I) \neq H_u$. For any vertex v define $K_v = \mathrm{Ker}(T_v - \lambda I)$. Then $f_\alpha(K_{s(\alpha)}) \subset (K_{r(\alpha)})$, for any edge $\alpha \in E$. In fact, for $x \in K_{s(\alpha)}$,

$$T_{r(\alpha)} f_\alpha x = f_\alpha T_{s(\alpha)} x = f_\alpha \lambda x = \lambda f_\alpha x.$$

Let g_α be the restriction of f_α to $K_{s(\alpha)}$. Then (K, g) is a non-trivial subrepresentation of (H, f).
The case (B): Define a subrepresentation (K, g) by $K_v = \mathrm{Ker}(T_v - \lambda_1 I)$ and $g_\alpha = f_\alpha|_{K_{s(\alpha)}}$. Since $K_{v_1} = H_{v_1}$ and $K_{v_2} = 0$. (K, g) is a non-trivial subrepresentation of (H, f). $\square$

Proposition 2.4. *Let* Γ *be a finite quiver with no oriented cycles and* (H, f) *a non-zero Hilbert representation of* Γ. *Then* (H, f) *is simple if and only if* (H, f) *is canonically simple.*

Proof. It is trivial that if (H, f) is canonically simple, then it is simple. Conversely assume that (H, f) is simple. Since Γ is a quiver with no oriented cycles, there is a sink v_0 in V. First consider the case that $H_{v_0} \neq 0$. Choose a non-zero vector $x \in H_{v_0}$. Define a representation (K, g) of Γ by $K_v = \mathbb{C}x$ if $v = v_0$ and $K_v = 0$ if $v \neq v_0$. Put $g_\alpha = 0$ for any $\alpha \in E$. Since v_0 is a sink, (K, g) is a non-zero subrepresentation of (H, f). Since (H, f) is simple, $(H, f) = (K, g)$. This means that (H, f) is canonically simple. Next consider the general case. Since Γ is a finite quiver with no oriented cycles, there exists a vertex v_1 in V such that $H_{v_1} \neq 0$ and $f_\alpha = 0$ for any edge $\alpha \in E$ with $s(\alpha) = v_1$. Choose a non-zero vector $y \in H_{v_1}$. Define a representation (L, h) of Γ by $L_v = \mathbb{C}y$ if $v = v_1$ and $L_v = 0$ if $v \neq v_1$. Put $h_\alpha = 0$ for any $\alpha \in E$. Then (L, h) is a non-zero subrepresentation of (H, f). Since (H, f) is simple, $(H, f) = (L, h)$. Therefore (H, f) is canonically simple. ☐

Proposition 2.5. *Let* L_2 *be 2-loop, that is,* L_2 *is a quiver with one vertex* $\{v\}$ *and two loops* $\{\alpha, \beta\}$. *Let* T *be a bounded operator on an infinite dimensional Hilbert space* H. *Let* (H, f) *be an infinite dimensional representation of* Γ *such that* $H_v = H$ *and* $f_\alpha = T, f_\beta = T^*$. *Then the following conditions are equivalent:*

1. (H, f) *is transitive .*
2. (H, f) *is simple.*
3. T *is irreducible, that is, the commutant* $\{T, T^*\}' = \mathbb{C}$.

Proof. Note that $End(H, f) = \{A \in B(H) \mid AT = TA,\ AT^* = T^*A\}$. Any subrepresentation is given by a subspace M of H such that $TM \subset M$ and $T^*M \subset M$. Let P be the projection of H onto M. Then this means that P commutes with T and T^*. Therefore these three condition (1), (2) and (3) are equivalent. ☐

3. HILBERT REPRESENTATIONS OF THE KRONECKER QUIVER

The Kronecker quiver K is a quiver with two vertices $\{1, 2\}$ and two paralleled arrows $\{\alpha, \beta\}$:

$$K : 1 \underset{\beta}{\overset{\alpha}{\rightrightarrows}} 2$$

A Hilbert representation (H, f) of the Kronecker quiver is given by two Hilbert spaces H_1, H_2 and two bounded operators $f_\alpha, f_\beta : H_1 \to H_2$.

The finite-dimensional indecomposable representations of the Kronecker quiver K was partially classified by Weierstrass and completed by Kronecker. Any finite-dimensional indecomposable representation is one of the following up to isomorphism:

(1) $H_1 = \mathbb{C}^n, H_2 = \mathbb{C}^n, f_\alpha = \lambda I_n + J_n,$ (Jordan block) $, \lambda \in \mathbb{C}, f_\beta = I_n, \quad n \geq 1.$

(2) $H_1 = \mathbb{C}^n, H_2 = \mathbb{C}^n, f_\alpha = I_n, f_\beta = \lambda I_n + J_n,$ (Jordan block) $, \lambda \in \mathbb{C}, \quad n \geq 1.$

(3) $H_1 = \mathbb{C}^{n+1}, H_2 = \mathbb{C}^n, f_\alpha = [I_n, 0], f_\beta = [0, I_n], n \geq 0.$

(4) $H_1 = \mathbb{C}^n, H_2 = \mathbb{C}^{n+1}, f_\alpha = \begin{bmatrix} I_n \\ 0 \end{bmatrix}, f_\beta = \begin{bmatrix} 0 \\ I_n \end{bmatrix}, n \geq 0.$

Moreover the case (2) is reduced to (1) if $\lambda \neq 0$. Among these cases, any representation (H, f) in (3),(4), n=1 of (1) or n = 1 of (2) is transitive and $End(H, f) = \mathbb{C}I$. Therefore it is interesting to find an infinite-dimensional indecomposable Hilbert representation (H, f) of the Kronecker quiver K and one with $End(H, f) = \mathbb{C}I$.

Example 8 Let K be the Kronecker quiver. Let S be the unilateral shift on $\ell^2(\mathbb{N})$. Define a Hilbert representation (H, f) of K by $H_1 = H_2 = \ell^2(\mathbb{N})$ and $f_\alpha = I, f_\beta = \lambda I + S, \ \lambda \in \mathbb{C}$. Then the Hilbert representation (H, f) of K is indecomposable and is not transitive.

Similarly, define a Hilbert representation (L, g) of K by $L_1 = L_2 = \ell^2(\mathbb{N})$ and $g_\alpha = I, g_\beta = \lambda I + S^*, \ \lambda \in \mathbb{C}$. Then the Hilbert representation (L, g) of K is indecomposable and is not transitive. These are infinite-dimensional analog of the case (1) and (2) of the finite-dimensional indecomposable representation of K.

We can replace the unilateral shift by a strongly irreducible operators in general. The following proposition shows that strongly irreducible operators are important to study Hilbert representations of quivers.

Proposition 3.1. *Let K be the Kronecker quiver. Let A be a bounded operator on a Hilbert space $\mathcal{H}$. Let (H, f) be a Hilbert representation defined by one of the following forms:*

(1) $H_1 = \mathcal{H}, H_2 = \mathcal{H}, f_\alpha = A, f_\beta = I.$

(2) $H_1 = \mathcal{H}, H_2 = \mathcal{H}, f_\alpha = I, f_\beta = A.$,

If A is strongly irreducible, then the Hilbert representation (H, f) of the Kronecker quiver K is indecomposable. The representation (H, f) is not transitive unless $\dim \mathcal{H} = 1$. *Conversely if the Hilbert representation (H, f) is indecomposable, then the operator A is strongly irreducible.*

Proof. Assume that A is strongly irreducible. Let $T = (T_1, T_2)$ be in $Idem(H, f)$. Then $T_2 I = I T_1$ and $T_2 A = A T_1$ in either case. Thus $T_1 A = A T_1$ and T_1 is an idempotent. Since A is strongly irreducible, $T_1 = 0$ or $T_1 = I$. Hence $T = 0$ or $T = I$, because $T_1 = T_2$. Therefore (H, f) is indecomposable. Furthermore suppose that $\dim \mathcal{H} \neq 1$. Since A is strongly irreducible, A is not a scalar operator. Then (A, A) is in $End(H, f)$ and is not a scalar. Therefore (H, f) is not transitive. Conversely, assume that (H, f) is indecomposable. Let $Q \in \mathcal{H}$ be an idempotent operator such that $QA = AQ$. Then $T = (Q, Q)$ is in $End(H, f)$. Since (H, f) is indecomposable, $Q = 0$ or $Q = I$. Hence A is strongly irreducible.

□

Lemma 3.2. *Let K be the Kronecker quiver. Let A and B be bounded operators on a Hilbert space $\mathcal{H}$. Let (H, f) be a Hilbert representation defined by $H_1 = \mathcal{H}, H_2 = \mathcal{H}, f_\alpha = A$ and $f_\beta = B$. If A is invertible, then there exists a bounded operator C on $\mathcal{H}$, such that (H, f) is isomorphic to a Hilbert representation (L, g) with $L_1 = \mathcal{H}, L_2 = \mathcal{H}, g_\alpha = I$ and $g_\beta = C$. Conversely if (H, f) is isomorphic to a Hilbert representation (L, g) with $g_\alpha = I$ and $g_\beta = C$, then A is invertible.*

Proof. Put $C = A^{-1}B$. Then $T = (T_1, T_2) := (A, I)$ gives a desired isomorphism of (H, f) onto (L, g). The converse is clear. □

Proposition 3.3. *Let K be the Kronecker quiver. Let A and B be bounded operators on a Hilbert space $\mathcal{H}$. Let (H, f) be a Hilbert representation defined by $H_1 = \mathcal{H}, H_2 = \mathcal{H}, f_\alpha = A$ and $f_\beta = B$. If $xA+yB$ is invertible for some scalars x and $y \neq 0$, then there exists a bounded operator T on $\mathcal{H}$, scalars $\lambda_0 \neq 0$ and λ_1 such that (H, f) is isomorphic to a Hilbert representation (L, g) with $L_1 = \mathcal{H}, L_2 = \mathcal{H}, g_\alpha = T$ and $g_\beta = \lambda_0 I + \lambda_1 T$.*

Proof. Put $T = (xA + yB)^{-1}A$. Then

$$(xA + yB)^{-1}B = (xA + yB)^{-1}(\frac{1}{y}(xA + yB) - \frac{x}{y}A) = \frac{1}{y}I - \frac{x}{y}T.$$

Therefore $T = (T_1, T_2) := (I, (xA+yB)^{-1})$ gives a desired isomorphism of (H, f) onto (L, g). □

We shall show that strongly irreducible operators are useful to construct indecomposable Hilbert representations of n-Kronecker quivers.

Theorem 3.4. *Let K_{n+1} be the $(n+1)$-Kronecker quiver, that is, K_{n+1} is a quiver with two vertex $\{1, 2\}$ and $n+1$ edges $\{\alpha_0, \dots, \alpha_n\}$ such that $s(\alpha_k) = 1$ and $r(\alpha_k) = 2$ for $k = 0, \dots, n$. Let T be a bounded operator on a Hilbert space H. Define a Hilbert representation $(M, f) = (M_T, f_T)$ of K_{n+1} by $M_1 = M_2 = H$ and $f_{\alpha_0} = \sum_{k=0}^n \lambda_k T^k$ for some scalars $\lambda_0 \neq 0, \lambda_1, \dots, \lambda_n$ and $f_{\alpha_k} = T^k$ for $k = 1, \dots, n$. Then the Hilbert representation (M, f) is indecomposable if and only if T is strongly irreducible. Moreover let S be another bounded operator. Then the corresponding Hilbert representations (M_T, f_T) and (M_S, f_S) are isomorphic if and only if T and S are similar.*

Proof. We shall show that

$$End(M, f) = \{(A, A) \in B(H)^2 \mid AT = TA\}$$

In fact, let $(A, B) \in B(H)^2$ be in $End(M, f)$. Then $BT^k = T^kA$ for $k = 1, \dots, n$ and $B(\sum_{k=0}^n \lambda_k T^k) = (\sum_{k=0}^n \lambda_k T^k)A$. Hence $\lambda_0 A = \lambda_0 B$.

Since $\lambda_0 \neq 0$, we have $A = B$. Then $AT = TA$. The converse is clear. Therefore

$$Idem(M, f) = \{(A, A) \in B(H)^2 \mid AT = TA, \quad A^2 = A\} = \{0, I\}$$

if and only if T is strongly irreducible. The latter half of the statement is similarly proved. □

Theorem 3.5. *Let K be the Kronecker quiver. Then there exist infinitely many infinite-dimensional indecomposable Hilbert representations of K which are relatively prime each other.*

Proof. Let S be the unilateral shift on $\ell^2(\mathbb{N})$. For each $\lambda \in \mathbb{C}$ define an indecomposable Hilbert representation (H^λ, f^λ) of K by $H_1^\lambda = H_2^\lambda = \ell^2(\mathbb{N})$ and $f_\alpha^\lambda = I, f_\beta^\lambda = \lambda I + S$. For $\lambda, \mu \in \mathbb{C}$, if $|\lambda - \mu| > 2$, then we shall show that (H^λ, f^λ) and (H^μ, f^μ) are relatively prime each other. On the contrary, assume that $Hom((H^\lambda, f^\lambda), (H^\mu, f^\mu)) \neq 0$. Then there exists a non-zero $T = (T_1, T_2) \in Hom((H^\lambda, f^\lambda), (H^\mu, f^\mu))$. Then

$$T_2 I = I T_1, \quad T_2(\lambda I + S) = (\mu I + S) T_1.$$

Hence $T_1 = T_2$ and $(\lambda - \mu) T_1 = S T_1 - T_1 S$. Then

$$2\|T_1\| < \|(\lambda - \mu) T_1\| = \|S T_1 - T_1 S\| \leq 2\|T_1\|.$$

This is a contradiction. Hence $Hom((H^\lambda, f^\lambda), (H^\mu, f^\mu)) = 0$. Similarly we have $Hom((H^\mu, f^\mu), (H^\lambda, f^\lambda)) = 0$. □

Since relatively prime Hilbert representations are not isomorphic, it is clear that there exist infinitely many non-isomorphic infinite-dimensional indecomposable Hilbert representations of K. We can easily say more.

Proposition 3.6. *Let K be the Kronecker quiver. Then there exist uncountably many non-isomorphic infinite-dimensional indecomposable Hilbert representations (H, f) of K.*

Proof. Since $\lambda I + S$ and $\mu I + S$ are not similar for $\lambda \neq \mu \in \mathbb{C}$, the Hilbert representations (H^λ, f^λ) and (H^μ, f^μ) of the Kronecker quiver K are not isomorphic. □

Example 9 Let K be the Kronecker quiver. Let S be the unilateral shift on $\ell^2(\mathbb{N})$. Define a Hilbert representation (H, f) of K by $H_1 = H_2 = \ell^2(\mathbb{N})$ and $f_\alpha = S, f_\beta = S^*$. Then the Hilbert representation (H, f) of K is not indecomposable, so that (H, f) of K is not simple. In fact, let $\{e_n \mid n \in \mathbb{N}\}$ be a canonical basis of $\ell^2(\mathbb{N})$. Define $L_1 = \{e_n \mid n \in \mathbb{N} \text{ is odd }\}$ and $L_2 = \{e_n \mid n \in \mathbb{N} \text{ is even }\}$. Consider the restrictions $g_\alpha = S|_{L_1}$ and $g_\beta = S^*|_{L_1}$. Then (L, g) is a non-trivial subrepresentation of (H, f). Similarly define $M_1 = \{e_n \mid n \in \mathbb{N} \text{ is even }\}$ and $M_2 = \{e_n \mid n \in \mathbb{N} \text{ is odd }\}$. Consider the restrictions $h_\alpha = S|_{M_1}$

and $h_\beta = S^*|_{M_1}$. Then (M, h) is also a non-trivial subrepresentation of (H, f) and $(H, f) = (L, g) \oplus (M, h)$.

In the following we shall construct infinite-dimensional indecomposable Hilbert representations of the Kronecker quiver which are transitive.

A perturbation of a weighted shift by a rank-one operator is crucially used to show being transitive.

Let H be a Hilbert space. Recall that for vectors $a, b \in H$, a rank-one operator $\theta_{a,b}$ is defined by $\theta_{a,b}(x) = (x|b)a, \quad x \in H$.

Theorem 3.7. *Let $K = (V, E, s, r)$ be the Kronecker quiver so that $V = \{1, 2\}$, $E = \{\alpha, \beta\}$ with $s(\alpha) = 1, s(\beta) = 1, r(\alpha) = 2$ and $r(\beta) = 2$,. Let S be the unilateral shift on $\mathcal{H} = \ell^2(\mathbb{N})$ with a canonical basis $\{e_1, e_2, ...\}$. For a bounded weight vector $\lambda = (\lambda_1, \lambda_2, ...) \in \ell^\infty$ we associate with a diagonal operator $D_\lambda = diag(\lambda_1, \lambda_2, ...)$, so that SD_λ is a weighted shift operator. We assume that $\lambda_i \neq \lambda_j$ if $i \neq j$. Take a vector $\overline{w} = (\overline{w_n})_n \in \ell^2(\mathbb{N})$ such that $w_n \neq 0$ for any $n \in \mathbb{N}$. Let (H, f) be a Hilbert representation such that $H_1 = H_2 = \mathcal{H} = \ell^2(\mathbb{N})$ and $f_\alpha = S, f_\beta = T := SD_\lambda + \theta_{e_1,\overline{w}}$. i.e., f_β is a perturbation of a weighted shift by a rank-one operator. Then the Hilbert representation (H, f) is transitive.*

Proof. We need an infinite matrix presentation of T:

$$T = SD_\lambda + \theta_{e_1,\overline{w}} = \begin{pmatrix} w_1 & w_2 & w_3 & ... \\ \lambda_1 & 0 & 0 & ... \\ 0 & \lambda_2 & 0 & ... \\ 0 & 0 & \lambda_3 & ... \\ 0 & 0 & 0 & ... \end{pmatrix}$$

Take any (ϕ, ψ) in $End(H, f)$. Write $\phi = (c_{ij})_{ij} \in B(\ell^2(\mathbb{N}))$ and

$$\psi = \begin{pmatrix} \alpha & \beta_1 & \beta_2 & \cdots \\ \gamma_1 & d_{11} & d_{12} & \cdots \\ \gamma_2 & d_{21} & d_{22} & \cdots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix} \in B(\ell^2(\mathbb{N})).$$

Then $S\phi = \psi S$ means that

$$\begin{pmatrix} 0 & 0 & 0 & \cdots \\ c_{11} & c_{12} & c_{13} & \cdots \\ c_{21} & c_{22} & c_{23} & \cdots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix} = \begin{pmatrix} \beta_1 & \beta_2 & \beta_3 & \cdots \\ d_{11} & d_{12} & d_{13} & \cdots \\ d_{21} & d_{22} & d_{23} & \cdots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix},$$

Hence we have $(c_{ij})_{ij} = (d_{ij})_{ij}$ and

$$\begin{pmatrix} \beta_1 & \beta_2 & \beta_3 & \cdots \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & \cdots \end{pmatrix}.$$

Therefore we have

$$\phi = \begin{pmatrix} c_{11} & c_{12} & ... \\ c_{21} & c_{22} & ... \\ \vdots & ... & ... \end{pmatrix}, \quad \text{and} \quad \psi = \begin{pmatrix} \alpha & 0 & 0 & ... \\ \gamma_1 & c_{11} & c_{12} & ... \\ \gamma_2 & c_{21} & c_{22} & ... \\ \vdots & ... & ... & ... \end{pmatrix}$$

Next we consider another compatibility condition $T\phi = \psi T$. Put $\widetilde{c_k} = \sum_{i=1}^{\infty} w_i c_{ik}$. Then we have

$$T\phi = \begin{pmatrix} w_1 & w_2 & w_3 & \cdots \\ \lambda_1 & 0 & 0 & \cdots \\ 0 & \lambda_2 & 0 & \cdots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix} \begin{pmatrix} c_{11} & c_{12} & c_{13} & \cdots \\ c_{21} & c_{22} & c_{23} & \cdots \\ c_{31} & c_{32} & c_{33} & \cdots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix}$$

$$= \begin{pmatrix} \widetilde{c_1} & \widetilde{c_2} & \widetilde{c_3} & \cdots \\ \lambda_1 c_{11} & \lambda_1 c_{12} & \lambda_1 c_{13} & \cdots \\ \lambda_2 c_{21} & \lambda_2 c_{22} & \lambda_2 c_{23} & \cdots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix}$$

We also have that

$$\psi T = \begin{pmatrix} \alpha & 0 & 0 & \cdots \\ \gamma_1 & c_{11} & c_{12} & \cdots \\ \gamma_2 & c_{21} & c_{22} & \cdots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix} \begin{pmatrix} w_1 & w_2 & w_3 & \cdots \\ \lambda_1 & 0 & 0 & \cdots \\ 0 & \lambda_2 & 0 & \cdots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix}$$

$$= \begin{pmatrix} \alpha w_1 & \alpha w_2 & \alpha w_3 & \cdots \\ \gamma_1 w_1 + c_{11}\lambda_1 & \gamma_1 w_2 + c_{12}\lambda_2 & \gamma_1 w_3 + c_{13}\lambda_3 \cdots & \\ \gamma_2 w_1 + c_{21}\lambda_1 & \gamma_2 w_2 + c_{22}\lambda_2 & \gamma_2 w_3 + c_{23}\lambda_3 \cdots & \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix}$$

Hence we have the following relations: $\alpha w_i = \widetilde{c_i} (i = 1, 2, 3. \cdots)$ and

$$\gamma_1 w_1 + c_{11}\lambda_1 = \lambda_1 c_{11},$$

$$\gamma_1 w_2 + c_{12}\lambda_2 = \lambda_1 c_{12},$$

$$\gamma_1 w_3 + c_{13}\lambda_3 = \lambda_1 c_{13}, \cdots .$$

Since $\gamma_1 w_1 + c_{11}\lambda_1 = \lambda_1 c_{11}$, we have $\gamma_1 w_1 = 0$. $w_1 \neq 0$ implies that $\gamma_1 = 0$. Therefore $c_{12}\lambda_2 = \lambda_1 c_{12}$. Since $\lambda_1 \neq \lambda_2$, we have $c_{12} = 0$. Similarly we have $c_{1j} = 0$ for $j \neq 1$.

Next look at the part including γ_2 parameter:

$$\gamma_2 w_1 + c_{21}\lambda_1 = \lambda_2 c_{21}, \gamma_2 w_2 + c_{22}\lambda_2 = \lambda_2 c_{22}, \gamma_2 w_3 + c_{23}\lambda_3 = \lambda_2 c_{23}, \cdots .$$

Since $\gamma_2 w_2 = 0$ and $w_2 \neq 0$, we have $\gamma_2 = 0$. By a similar argument as above, we have $c_{2j} = 0$ for $j \neq 2$. In the same way, we have $\gamma_i = 0$ for any i and $c_{ij} = 0$ for $i \neq j$. Therefore we have that

$$\psi = \begin{pmatrix} \alpha & 0 & 0 & \cdots \\ 0 & c_{11} & 0 & \cdots \\ 0 & 0 & c_{22} & \cdots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix}, \quad \text{and} \quad \phi = \begin{pmatrix} c_{11} & 0 & 0 & \cdots \\ 0 & c_{22} & 0 & \cdots \\ 0 & 0 & c_{33} & \cdots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix}.$$

Since $\widetilde{c_1} = \alpha w_1$ and $\widetilde{c_1} = w_1 c_{11} + w_2 c_{21} + w_3 c_{31} + \cdots = w_1 c_{11}$, we have $\alpha = c_{11}$, because $w_1 \neq 0$. Similarly the equations $\widetilde{c_2} = \alpha w_2$, $\widetilde{c_3} = \alpha w_3, \cdots$ and $w_i \neq 0$, we have $\alpha = c_{ii}$ for any i. Hence $(\phi, \psi) = (\alpha I, \alpha I)$. Thus (H, f) is transitive. □

Next we shall construct a transitive Hilbert representation of the Kronecker quiver in another method. It is a modification of an unbounded operator which provides a transitive lattice by Harrison,Radjavi and Rosenthal [HRR].

Let A and B be bounded operators on a Hilbert space $\mathcal{H}$. Let $(H^{(A,B)}, f^{(A,B)})$ be a Hilbert representation of the Kronecker quiver K defined by $H_1 = \mathcal{H}, H_2 = \mathcal{H}, f_\alpha = A$ and $f_\beta = B$. If A is invertible, then $(H^{(A,B)}, f^{(A,B)})$ is isomorphic to $(H^{(I,A^{-1}B)}, f^{(I,A^{-1}B)})$. Even if A does not have a bounded inverse, $A^{-1}B$ can be an unbounded operator. Hence if an unbounded operator C is formally written by $C =$ "$A^{-1}B$", then we might replace $(H^{(I,C)}, f^{(I,C)})$ by $(H^{(A,B)}, f^{(A,B)})$ to keep it in the category of bounded operators. We shall adapt the idea to an unbounded operator C which gives a transitive lattice in [HRR]. We also modify it a little bit to make a calculation of $End(H, f)$ easier.

Theorem 3.8. *Let K be the Kronecker quiver and $\mathcal{H} = \ell^2(\mathbb{Z})$. Fix a positive constant $\lambda > 1$. Consider two weight sequences $a = (a(n))_{n\in\mathbb{Z}}$ and $b = (b(n))_{n\in\mathbb{Z}}$ by*

$$a(n) = \begin{cases} e^{-\lambda^n}, & (n \geq 1, n \textit{ is even }) \\ 1, & (\textit{otherwise}), \end{cases} \qquad b(n) = \begin{cases} e^{-\lambda^n}, & (n \geq 1, n \textit{ is odd }) \\ 1, & (\textit{otherwise}). \end{cases}$$

Let D_a be a diagonal operator with $a = (a(n))_n$ as diagonal coefficients. and D_b be a diagonal operator with $b = (b(n))_n$ as diagonal coefficients. Let U be the bilateral forward shift. Put $A = D_a$ and $B = UD_b$, so that A is a positive operator and B is a weighted forward shift operator. Define a Hilbert representation (H^λ, f^λ) of the Kronecker quiver K by $H_0^\lambda = \mathcal{H}, H_1^\lambda = \mathcal{H}$, $f_\alpha^\lambda = A$ and $f_\beta^\lambda = B$. Then the Hilbert representation (H^λ, f^λ) of K is transitive and is not isomorphic to any of the Hilbert representation in Theorem 3.7 constructed by a perturbation of a weighted shift by a rank-one operator.

Proof. Let $\{e_n\}_{n\in\mathbb{Z}}$ be the canonical basis of $\mathcal{H} = \ell^2(\mathbb{Z})$.

Let $T = (T_1, T_2)$ be in $End(H^\lambda, f^\lambda)$. Then $T_2 A = AT_1$ and $T_2 B = BT_1$, that is, $T_2 D_a = D_a T_1$ and $T_2 U D_b = U D_b T_1$. Since $(T_2 D_a e_n \mid$

$e_m) = (D_a T_1 e_n \mid e_m)$, we have $a(n)(T_2 e_n \mid e_m) = a(m)(T_1 e_n \mid e_m)$. Hence

$$(T_1 e_n \mid e_m) = \frac{a(n)}{a(m)}(T_2 e_n \mid e_m).$$

Since $(T_2 U D_b e_n \mid e_m) = (U D_b T_1 e_n \mid e_m)$, we have $b(n)(T_2 e_{n+1} \mid e_m) = (T_1 e_n \mid b(m-1)e_{m-1})$. Replacing m by $m+1$, we obtain

$$(T_1 e_n \mid e_m) = \frac{b(n)}{b(m)}(T_2 e_{n+1} \mid e_{m+1}).$$

Combining these equations, we have $\frac{a(n)}{a(m)}(T_2 e_n \mid e_m) = \frac{b(n)}{b(m)}(T_2 e_{n+1} \mid e_{m+1})$. Put $w_m = \frac{b(m)}{a(m)}$. Then $w_m = e^{(-\lambda)^m}$ for $m \geq 1$.

$$(T_2 e_{n+1} \mid e_{m+1}) = \frac{w_m}{w_n}(T_2 e_n \mid e_m).$$

Since $m-n$ component $t_{m,n}$ of T_2 is given by $t_{m,n} = (T_2 e_n \mid e_m)$, $t_{m+1,n+1} = \frac{w_m}{w_n} t_{m,n}$. Putting $m = n$, we have $t_{n+1,n+1} = t_{n,n}$. This shows that the diagonal of T_2 is a constant. We shall show that T_2 is a scalar operator. Suppose that T_2 were not a scalar operator. Then there exist integers $m \neq n$ such that $t_{m,n} \neq 0$. For any integer $k \geq 1$, we have

$$t_{m+k,n+k} = \frac{w_m w_{m+1} \cdots w_{m+k-1}}{w_n w_{n+1} \cdots w_{n+k-1}} t_{m,n}.$$

Define $c_k(m,n)$ by $\frac{w_m w_{m+1} \cdots w_{m+k-1}}{w_n w_{n+1} \cdots w_{n+k-1}} = e^{c_k(m,n)}$. First consider the case that $m \geq 1$ and $n \geq 1$. Then we have

$$\begin{aligned} c_k(m,n) &= ((-\lambda)^m + (-\lambda)^{m+1} + \cdots + (-\lambda)^{m+k-1}) \\ &\quad - ((-\lambda)^n + (-\lambda)^{n+1} + \cdots + (-\lambda)^{n+k-1}) \\ &= \frac{(-\lambda)^m(1-(-\lambda)^k)}{1+\lambda} - \frac{(-\lambda)^n(1-(-\lambda)^k)}{1+\lambda}. \end{aligned}$$

Since $\limsup_k c_k(m,n) = \infty$, we have $\limsup_k t_{m+k,n+k} = \infty$. This contradicts to the boundedness of T_2. The other cases are similarly proved. Consequently $T_2 = \alpha I$ for some constant α.

Since $(T_1 e_n \mid e_m) = \frac{a(n)}{a(m)}(T_2 e_n \mid e_m)$, we have $(T_1 e_n \mid e_m) = 0$ for $m \neq n$ and $(T_1 e_n \mid e_n) = \alpha$ for any $n \in \mathbb{Z}$. Hence $T_1 = T_2 = \alpha I$. This shows that the Hilbert representation (H^λ, f^λ) is transitive.

The Hilbert representation (H^λ, f^λ) is not isomorphic to any of the Hilbert representation (H, f) in 3.7 constructed by a perturbation of a weighted shift by a rank-one operator. In fact the image of f_α is closed but the image of f_α^λ is not closed.

□

A little more careful calculation shows that (H^λ, f^λ) and (H^μ, f^μ) are not isomorphic if $\lambda \neq \mu$, $\lambda > 1$ and $\mu > 1$ as follows:

Theorem 3.9. *Let K be the Kronecker quiver. Then there exist continuously many non-isomorphic Hilbert representations $(H^{\lambda}, f^{\lambda})$ of the Kronecker quiver K which are transitive.*

Proof. Let λ and μ be positive constants such that $\lambda \neq \mu,\ \lambda > 1$ and $\mu > 1$. It is enough to show that $Hom((H^{\lambda}, f^{\lambda}), (H^{\mu}, f^{\mu})) = 0$. We shall write $A^{\mu}, B^{\mu}, a^{\mu}$ and b^{μ} for (H^{μ}, f^{μ}). Take any homomorphism $(T_1, T_2) \in Hom((H^{\lambda}, f^{\lambda}), (H^{\mu}, f^{\mu}))$. Then we have $T_2A^{\lambda} = A^{\mu}T_1$ and $T_2B^{\lambda} = B^{\mu}T_1$. Since $(T_2A^{\lambda}e_n \mid e_m) = (A^{\mu}T_1e_n \mid e_m)$, we have $a^{\lambda}(n)(T_2e_n \mid e_m) = (T_1e_n \mid a^{\mu}(m)e_m)$. Hence we obtain that

$$(T_1e_n \mid e_m) = \frac{a^{\lambda}(n)}{a^{\mu}(m)}(T_2e_n \mid e_m)$$

Next consider $(T_2B^{\lambda}e_n \mid e_m) = (B^{\mu}T_1e_n \mid e_m)$. Since $(T_2UD_b^{\lambda}e_n \mid e_m) = (T_1e_n \mid D_b^{\mu}U^*e_m)$, we have that $b^{\lambda}(n)(T_2e_{n+1} \mid e_m) = b^{\mu}(m-1)(T_1e_n \mid e_{m-1})$. Replacing m by $m+1$, we obtain

$$(T_1e_n \mid e_m) = \frac{b^{\lambda}(n)}{b^{\mu}(m)}(T_2e_{n+1} \mid e_{m+1}).$$

Combining these equations, we have

$$(T_2e_{n+1} \mid e_{m+1}) = \frac{a^{\lambda}(n)}{a^{\mu}(m)}\frac{b^{\mu}(m)}{b^{\lambda}(n)}(T_2e_n \mid e_m)$$

.

We put $w_m^{\mu} = \frac{b^{\mu}(m)}{a^{\mu}(m)}$. Then

$$(T_2e_{n+1} \mid e_{m+1}) = \frac{w_m^{\mu}}{w_n^{\lambda}}(T_2e_n \mid e_m).$$

For any integer $k \geq 1$, we have

$$(T_2e_{n+k} \mid e_{m+k}) = \frac{w_m^{\mu} \cdots w_{m+k-1}^{\mu}}{w_n^{\lambda} \cdots w_{n+k-1}^{\lambda}}(T_2e_n \mid e_m)$$

Define $c_k(m,n)$ by

$$\frac{w_m^{\mu} \cdots w_{m+k-1}^{\mu}}{w_n^{\lambda} \cdots w_{n+k-1}^{\lambda}} = e^{c_k(m,n)}.$$

Then we have

$$c_k(m,n) = ((-\mu)^m + \cdots (-\mu)^{m+k-1}) - ((-\lambda)^n + \cdots (-\lambda)^{n+k-1})$$

Since $\mu \neq \lambda$, we may and do assume that $1 < \lambda < \mu$. We shall show that $T_2 = 0$. On the contrary, assume that $T_2 \neq 0$. Then there exist integers m, n such that $(T_2e_n \mid e_m) \neq 0$. We can show that $\limsup_k c_k(m,n) = \infty$. For example, if $m \geq 1$ and $n \geq 1$, then

$$c_k(m,n) = \frac{(-\mu)^m(1-(-\mu)^k)}{1+\mu}\{1 - \frac{(-\lambda)^n(1+\mu)(1-(-\lambda)^k)}{(-\mu)^m(1+\lambda)(1-(-\mu)^k)}\}$$

Hence $\limsup_k c_k(m,n) = \infty$. The rest cases are similarly proved. But this contradicts that T_2 is bounded. Therefore $T_2 = 0$. Since $(T_1 e_n \mid e_m) = \frac{a^\lambda(n)}{a^\mu(m)}(T_2 e_n \mid e_m)$, we also have T_1=0. This shows that $T = (T_1, T_2) = 0$. Hence we have that $Hom((H^\lambda, f^\lambda), (H^\mu, f^\mu)) = 0$. Therefore (H^λ, f^λ) is not isomorphic to (H^μ, f^μ). $\square$

4. Difference between purely algebraic version and Hilbert space version

There exist subtle difference among purely algebraic infinite-dimensional representations of quivers, infinite-dimensional Banach (space) representations of quivers and infinite-dimensional Hilbert (space) representations of quivers. We also note that the analytic aspect of Hardy space is quite important in our setting.

We shall use the following elementary fact: Let A be a unital algebra and $L(A)$ be the set of linear operators on A. . Let $\lambda : A \to L(A)$ be the left multiplication such that $\lambda_a x = ax$ for $a, x \in A$. Similarly let $\rho : A \to L(A)$ be the right multiplication such that $\rho_a x = xa$ for $a, x \in A$. Then the commutant

$$\lambda(A)' := \{T \in L(A) \mid T\lambda_a = \lambda_a T \text{ for any } a \in A\}$$

is exactly $\rho(A)$. In fact, let $T \in \lambda(A)'$. Put $b = T(1) \in A$. Then $T\lambda_a(1) = \lambda_a T(1)$ for any $a \in A$. Hence $T(a) = ab = \rho_b(a)$. Therefore $T = \rho_b$.

Definition. Let $\Gamma = (V, E, s, r)$ be a finite quiver. We say that (K, f) is a *Banach representation* of Γ if $K = (K_v)_{v \in V}$ is a family of Banach spaces and $f = (f_\alpha)_{\alpha \in E}$ is a family of bounded linear operators such that $f_\alpha : K_{s(\alpha)} \to K_{r(\alpha)}$ for $\alpha \in E$. A Banach representation (K, f) of Γ is called *decomposable* if (K, f) is isomorphic to a direct sum of two non-zero Banach representations. A non-zero Banach representation (H, f) of Γ is said to be *indecomposable* if it is not decomposable. The other notions are similarly defined for Banach representations. A Banach representation (H, f) is indecomposable if and only if any endomorphism of (H, f) which is idempotent is 0 or I.

Example 10. Let L_1 be one-loop quiver, that is, L_1 is a quiver with one vertex $\{1\}$ and 1-loop $\{\alpha\}$ such that $s(\alpha) = r(\alpha) = 1$. Consider a purely algebraic group algebra $V_1 := \mathbb{C}[\mathbb{Z}]$, the reduced group C^*-algebra $K_1 := C^*_r(\mathbb{Z})$, the group von Neumann algebra $W^*(\mathbb{Z})$ and a Hilbert space $H_1 = \ell^2(\mathbb{Z})$. We identify V_1 with the algebra of finite Laurant polynomials. As a set we have inclusions under a certain identification:

$$V_1 = \mathbb{C}[\mathbb{Z}] \subset K_1 = C^*_r(\mathbb{Z}) \subset W^*(\mathbb{Z}) \subset \ell^2(\mathbb{Z}) = H_1.$$

Moreover V_1 is a dense subset of H_1 with respect to the topology of ℓ^2 norm of H_1. Define a purely algebraic representation (V, T) of L_1 by

$V_1 = \mathbb{C}[\mathbb{Z}]$ and the multiplication operator T_α by z. That is, $T_\alpha h(z) = zh(z)$ for a finite Laurant polynomial $h(z) = \sum_n a_n z^n \in \mathbb{C}[\mathbb{Z}]$. Similarly we can define a Banach space representation (K, S) of L_1 by $K_1 = C_r^*(\mathbb{Z}) \cong C(\mathbb{T})$ and the multiplication operator S_α by z.

Since

$$End(V, T) \cong \mathbb{C}[\mathbb{Z}] \subset C_r^*(\mathbb{Z}) \cong C(\mathbb{T})$$

and $C(\mathbb{T})$ have no non-trivial idempotents, the purely algebraic representation (V, T) and the Banach representation (K, S) is indecomposable. On the other hand, the multiplication operator U_α by z on $H_1 = \ell^2(\mathbb{Z}) \cong L^2(\mathbb{T})$ gives a Hilbert representation (H, U). Since U is a unitary, for any operator A, $AU_\alpha = U_\alpha A$ implies $AU_\alpha^* = U_\alpha^* A$. Therefore

$$End(H, U) \cong \{A \in B(\ell^2(\mathbb{Z})) \mid AU_\alpha = U_\alpha A\} \cong L^\infty(\mathbb{T}).$$

because $L^\infty(\mathbb{T})$ is maximal abelian in $B(L^2(\mathbb{T})) \cong B(\ell^2(\mathbb{Z}))$. Since $L^\infty(\mathbb{T})$ has many non-trivial idempotents, the Hilbert representation (H, U) is *not* indecomposable. Therefore the completion by the L^2-norm changes the indecomposability but the the completion by the sup-norm does not change the indecomposability. The example suggests that proving indecomposability for Hilbert representations is sometimes more difficult than proving indecomposability for purely algebraic representations.

We shall extend the above example to the n-loop quiver.

Proposition 4.1. *Let L_n be the n-loop quiver, that is, L_n is a quiver with one vertex $\{1\}$ and n-loops $\{\alpha_k \; | k = 1, \dots, n\}$ with $s(\alpha_k) = r(\alpha_k) = 1$ for $k = 1, \dots, n$. Let F_n be the (non-abelian) free group of n-generators $\{a_1, \dots, a_n\}$. Consider the purely algebraic group algebra $V_1 := \mathbb{C}[F_n]$, the reduced group C^*-algebra $K_1 =: C_r^*(F_n)$, the group von Neumann algebra $W^*(F_n)$ and a Hilbert space $H_1 = \ell^2(F_n)$. Let $\{\delta_g \mid g \in F_n\}$ be a standard basis of $H_1 = \ell^2(F_n)$ such that $\delta_g(h) = 1$ if $g = h$ and $\delta_g(h) = 0$ if $g \neq h$. Let $\lambda : F_n \to B(\ell^2(F_n))$ be the left regular representation, that is, $\lambda_s(\delta_g) = \delta_{sg}$. The left regular representation defines a Hilbert representation (H, f) by $H_1 = \ell^2(F_n)$ and $f_{\alpha_k} = \lambda_{a_k}$ for $k = 1, \dots, n$. Define a Banach representation (K, U) by its restriction to $C_r^*(F_n)$, that is, $K_1 = C_r^*(F_n)$ and $U_{\alpha_k} = \lambda_{a_k}|_{K_1}$ for $k = 1, \dots, n$. We also consider a purely algebraic representation (V, T) by representation its restriction to $\mathbb{C}[F_n]$, that is, $V_1 = \mathbb{C}[F_n]$ and $T_{\alpha_k} = \lambda_{a_k}|_{V_1}$ for $k = 1, \dots, n$. Then the purely algebraic representation (V, T) and Banach representation (K, U) are indecomposable but the Hilbert representation (H, f) is not indecomposable.*

Proof. The reduced group C^*-algebra $C_r^*(F_n)$ is the C^*-algebra generated by $\{\lambda_s \mid s \in F_n\}$ and has no non-trivial idempotents. The group von Neumann algebra $W^*(F_n)$ is the von Neumann algebra generated by $\{\lambda_s \mid s \in F_n\}$ and has many non-trivial idempotents. The purely algebraic group algebra $\mathbb{C}[F_n]$ is dense in $C_r^*(F_n)$ with respect to operator norm and dense in $W^*(F_n)$ with respect to weak operator topology. Moreover $\mathbb{C}[F_n]$ is dense in $\ell^2(F_n)$ with respect to ℓ^2 norm. Define an embedding $\eta : W^*(F_n) \to \ell^2(F_n)$ by $\eta(T) = T\delta_1$. In this sense we have inclusions as set:

$$\mathbb{C}[F_n] \subset C_r^*(F_n) \subset W^*(F_n) \subset \ell^2(F_n).$$

Since

$$End(H, f) \cong \{A \in B(\ell^2(\mathbb{Z})) \mid AB = BA \text{ for any } B \in W^*(F_n)\}$$

is the von Neumann algebra generated by the *right* regular representation and isomorphic to $W^*(F_n)$, $End(H, f)$ has many non-trivial idempotent. Hence the Hilbert representation (H, f) is *not* indecomposable. On the other hand,

$$End(K, U) = C^*(\{\rho_s \mid s \in F_n\}) \cong C_r^*(F_n).$$

Since $C_r^*(F_n)$ has no non-trivial projections, $C_r^*(F_n)$ has no non-trivial idempotents. Hence the Banach representation (K, U) is indecomposable. Since

$$End(V, T) \cong \mathbb{C}[F_n] \subset C_r^*(F_n)$$

have no idempotents, the purely algebraic representation (V, T) is also indecomposable. □

The same phenomenon occurs for $(n + 1)$-Kronecker quiver by a similar argument.

Proposition 4.2. *Let K_{n+1} be the $n + 1$-Kronecker quiver , that is, K_{n+1} is a quiver with two vertex $\{1, 2\}$ and $n + 1$ edges $\{\alpha_1, \dots, \alpha_{n+1}\}$ such that $s(\alpha_k) = 1$ and $r(\alpha_k) = 2$ for $k = 1, \dots, n + 1$. Let F_n be the (non-abelian) free group of n-generators $\{a_1, \dots, a_n\}$. Define a Hilbert representation (H, f) of K_{n+1} by $H_1 = H_2 = \ell^2(F_n)$ and $f_{\alpha_k} = \lambda_{a_k}$ for $k = 1, \dots, n$ and $f_{\alpha_{n+1}} = I$. We also define a Banach representation (K, U) and a purely algebraic representation (V, T) of K_{n+1} by the restriction to $C_r^*(F_n)$ and $\mathbb{C}[F_n]$ respectively: Let $K_1 = K_2 = C_r^*(F_n)$ and $U_{\alpha_k} = \lambda_{a_k}|_{K_1}$ for $k = 1, \dots, n + 1$. We also put $V_1 = V_2 = \mathbb{C}[F_n]$ and $T_{\alpha_k} = \lambda_{a_k}|_{V_1}$ for $k = 1, \dots, n+1$. Then the purely algebraic representation (V, T) and Banach representation (K, U) are indecomposable but the Hilbert representation (H, f) is not indecomposable.*

Proof. Since $End(H, f) \cong \{(A, A) \in B(\ell^2(\mathbb{Z}))^2 \mid AB = BA \text{ for any } B \in W^*(F_n)\}$ is isomorphic to $W^*(F_n)$ and $End(H, f)$ has many non-trivial idempotents, the Hilbert representation (H, f) is *not* indecomposable. On the other hand,

$$End(K, U) \cong \{(A, A) \in (C_r^*[F_n])^2 \mid A \in C_r^*[F_n]\}$$

and

$$End(V, T) \cong \{(A, A) \in \mathbb{C}[F_n]^2 \mid A \in \mathbb{C}[F_n]\},$$

have no non-trivial idempotents, (K,U) and (V,T) are indecomposable. □

Example 11. Let L_1 be one-loop quiver, that is, L_1 is a quiver with one vertex $\{1\}$ and 1-loop $\{\alpha\}$ such that $s(\alpha) = r(\alpha) = 1$. Consider two infinite-dimensional spaces the polynomial ring $\mathbb{C}[z]$ and the Hardy space $H^2(\mathbb{T})$. Then $\mathbb{C}[z]$ is dense in $H^2(\mathbb{T})$ with respect to the Hilbert space norm.

Define a purely algebraic representation (V,T) of L_1 by $V_1 = \mathbb{C}[z]$ and the multiplication operator T_α by z. That is, $T_\alpha h(z) = zh(z)$ for a polynomial $h(z) = \sum_n a_n z^n$. Since $End(V,T) \cong \mathbb{C}[z]$ have no idempotents, the purely algebraic representation (V,T) is indecomposable.

Next we define a Hilbert representation (H,S) by $H_1 = H^2(\mathbb{T})$ and $S_\alpha = T_z$ the Toeplitz operator with the symbol z. Then $S_\alpha = T_z$ is the multiplication operator by z on $H^2(\mathbb{T})$ and is identified with the unilateral shift. Then

$$\begin{aligned} End(H,S) &\cong \{A \in B(H^2(\mathbb{T})) \mid AT_z = T_z A\} \\ &= \{T_\phi \in B(H^2(\mathbb{T})) \mid \phi \in H^\infty(\mathbb{T})\} \end{aligned}$$

is the algebra of analytic Toeplitz operators and isomorphic to $H^\infty(\mathbb{T})$. By the F. and M. Riesz Theorem, if $f \in H^2(\mathbb{T})$ has the zero set of positive measure, then $f = 0$. Since $H^\infty(\mathbb{T}) = H^2(\mathbb{T}) \cap L^\infty(\mathbb{T})$, $H^\infty(\mathbb{T})$ has no non-trivial idempotents. Thus there exists no non-trivial idempotents which commutes with T_z and Hilbert space (H,S) is indecomposable. In this sense, the analytical aspect of Hardy space is quite important in our setting.

Any subrepresentation of the purely algebraic representation (V,T) is given by the restriction to an ideal $J = p(z)\mathbb{C}[z]$ for some polynomial $p(z)$. Any subrepresentation of the Hilbert representation (H,S) is given by an invariant subspace of the shift operator T_z. Beurling theorem shows that any subrepresentation of (H,S) is given by the restriction to an invariant subspace $M = \varphi H^2(\mathbb{T})$ for some inner function φ. For example, if an ideal J is defined by

$$J = \{f(z) \in \mathbb{C}[z] \mid f(\lambda_1) = \cdots = f(\lambda_n) = 0\}$$

for some distinct numbers $\lambda_1, \ldots \lambda_n \in \mathbb{C}$, then the corresponding polynomial $p(z)$ is given by $p(z) = (z - \lambda_1)\ldots(z - \lambda_n)$. The case of Hardy space is much more analytic. We shall identify $H^2(\mathbb{T})$ with a subspace $H^2(\mathbb{D})$ of analytic functions on the open unit disc $\mathbb{D}$. If an invariant subspace M is defined by

$$M = \{f \in H^2(\mathbb{D}) \mid \ f(\lambda_1) = \cdots = f(\lambda_n) = 0\}$$

for some distinct numbers $\lambda_1, \dots \lambda_n \in \mathbb{D}$, then the corresponding inner function φ is given by a finite Blaschke product

$$\varphi(z) = \frac{(z - \lambda_1)}{1 - \overline{\lambda_1} z} \cdots \frac{(z - \lambda_n)}{1 - \overline{\lambda_n} z}.$$

Here we cannot use the notion of degree like polynomials and we must manage to treat orthogonality to find such an inner function φ.

5. Hilbert representations and relative position of subspaces

We studied relative position of subspaces of a Hilbert space in [EW1], [EW2] and Hilbert representations of quivers in [EW3]. In this section we shall describe a relation between them, which is similar to purely algebraic situation and easy but quite suggestive. Therefore we shall describe it here.

Let H be a Hilbert space and $E_1, \dots E_n$ be n-subspaces in H. Then we say that $\mathcal{S} = (H; E_1, \dots, E_n)$ is a system of n-subspaces in H. Let $\mathcal{T} = (K; F_1, \dots, F_n)$ be another system of n-subspaces in a Hilbert space K. Then $\varphi : \mathcal{S} \to \mathcal{T}$ is called a homomorphism if $\varphi : H \to K$ is a bounded linear operator satisfying that $\varphi(E_i) \subset F_i$ for $i = 1, \dots, n$. And $\varphi : \mathcal{S} \to \mathcal{T}$ is called an isomorphism if $\varphi : H \to K$ is an invertible (i.e., bounded bijective) linear operator satisfying that $\varphi(E_i) = F_i$ for $i = 1, \dots, n$. We say that systems $\mathcal{S}$ and $\mathcal{T}$ are *isomorphic* if there is an isomorphism $\varphi : \mathcal{S} \to \mathcal{T}$. This means that the relative positions of n subspaces $(E_1, \dots, E_n)$ in H and $(F_1, \dots, F_n)$ in K are same under disregarding angles. We denote by $Hom(\mathcal{S}, \mathcal{T})$ the set of homomorphisms of $\mathcal{S}$ to $\mathcal{T}$ and $End(\mathcal{S}) := Hom(\mathcal{S}, \mathcal{S})$ the set of endomorphisms on $\mathcal{S}$. A system $\mathcal{S} = (H; E_1, \dots, E_n)$ of n subspaces is called *decomposable* if the system $\mathcal{S}$ is isomorphic to a direct sum of two non-zero systems. A non-zero system $\mathcal{S} = (H; E_1, \cdots, E_n)$ of n-subspaces is said to be *indecomposable* if it is not decomposable.

We recall that strongly irreducible operators A play a crucial role to construct indecomposable systems of four subspaces. Moreover the commutant $\{A\}'$ corresponds to the endomorphism ring.

Theorem 5.1 ([EW1]). *For any single operator $A \in B(K)$ on a Hilbert space K, let $\mathcal{S}_A = (H; E_1, E_2, E_3, E_4)$ be the associated operator system such that $H = K \oplus K$ and*

$$E_1 = K \oplus 0, E_2 = 0 \oplus K, E_3 = \{(x, Ax); x \in K\}, E_4 = \{(y, y); y \in K\}.$$

Then

$$End(\mathcal{S}_A) = \{T \oplus T \in B(H); T \in B(K),\ AT = TA\}$$

is isomorphic to the commutant $\{A\}$. The associated system $\mathcal{S}_A$ of four subspaces is indecomposable if and only if A is strongly irreducible. Moreover for any operators $A, B \in B(K)$ on a Hilbert space K, the *associated systems $\mathcal{S}_A$ and $\mathcal{S}_B$ are isomorphic if and only if A and B are similar.*

Example 12. We shall apply the above theorem to the famous facts on weighted shift operators and analytic function theory, see A. Shields [Sh]. Let $a = (a_n)_{n\in\mathbb{N}}$ be a bounded sequence and W_a be the associated weighted unilateral shift. Then the associated system $\mathcal{S}_{W_a}$ of four subspaces is indecomposable if and only if W_a is strongly irreducible if and only if $a_n \neq 0$ for any $n \in \mathbb{N}$. Let $b = (b_n)_{n\in\mathbb{N}}$ be another bounded sequence. Then the associated system $\mathcal{S}_{W_a}$ and $\mathcal{S}_{W_b}$ are isomorphic if and only if W_a and W_b are similar if and only if there exist positive constants C_1 and C_2 such that for any $n \in \mathbb{N}$

$$0 < C_1 \leq \frac{|a_1 \dots a_n|}{|b_1 \dots b_n|} \leq C_2.$$

Moreover $End(\mathcal{S}_{W_a})$ is isomorphic to the commutant $\{W_a\}'$, which is isomorphic to the "analytic" algebra $H^\infty(\beta)$ in the sense of [Sh], that is, the class of formal power series ϕ such that $\phi H^2(\beta) \subset H^2(\beta)$, where $\beta_0 = 1$, $\beta_n = a_0 a_1 \dots a_{n-1}$ and

$$H^2(\beta) = \{f | f(z) = \sum_{n=0}^{\infty} c_n z^n,\ \sum_{n=0}^{\infty} |c_n \beta_n|^2 < \infty\}.$$

It is easy but fundamental to see that the study of relative positions of subspaces is reduced to the study of Hilbert representations of quivers. In particular the indecomposabilty is preserved: Let $\mathcal{S} = (H; E_1, \dots, E_n)$ be a system of n-subspaces in a Hilbert space H. Let $R_n = (V, E, s, r)$ be a n subspace quiver such that $V = \{1, 2, \dots, n, n+1\}$ and $E = \{\alpha_k \mid k = 1, \dots, n\}$ with $s(\alpha_k) = k$ and $r(\alpha_k) = n + 1$ for $k = 1, \dots, n$. Then there exists a Hilbert representation (K, f) of R_n such that $K_k = E_k$, $K_{n+1} = H$ and $f_{\alpha_k} : E_k \to H$ is an inclusion for $k = 1, \dots, n$. Then there exists an algebra isomorphism

$$\theta : End(\mathcal{S}) \to End(K, f)$$

such that $\theta(\varphi) = (\varphi | K_k)_{k\in V}$ for $\varphi \in End(\mathcal{S})$. Therefore it is clear that the system $\mathcal{S}$ of n subspaces is indecomposable (resp. transitive) if and only if the corresponding Hilbert representation (K, f) is indecomposable (resp. transitive).

We shall show a converse in a sense as same as the purely algebraic version.

Lemma 5.2. *Let $\Gamma = (V, E, s, r)$ be a finite quiver without self-loops such that $V = \{v_1, \dots, v_n\}$ and $E = \{\alpha_1, \dots, \alpha_m\}$. Let (K, f) be a Hilbert representation of Γ. Then there exists a system $\mathcal{S} = (H; E_1, \dots, E_{n+m})$ of $n + m$-subspaces such that $End(K, f) \cong End(\mathcal{S})$.*

Proof. Let $H = K_{v_1} \oplus K_{v_2} \oplus \cdots \oplus K_{v_n}$. Define $E_i = 0 \oplus \cdots \oplus K_{v_i} \oplus \cdots \oplus 0$ for $i = 1, 2, \ldots, n$. For $j = 1, \ldots, m$, consider

$$f_{\alpha_j} : K_{s(\alpha_j)} \to K_{r(\alpha_j)}.$$

Define

$$\begin{aligned}&E_{n+j} = \text{"}graph\ f_{\alpha_j}\text{"} := \{x = (x_k)_k \in H \mid \\ &x = (0, \ldots, 0, z, \ldots, f_{\alpha_j}(z), 0 \ldots, 0)\ z \in K_{s(\alpha_j)}, f_{\alpha_j}(z) \in K_{r(\alpha_j)}\}.\end{aligned}$$

Then $\mathcal{S} := (H; E_1, \ldots, E_{n+m})$ is a system of $n + m$-subspaces.

For $T = (T_{v_k})_k \in End(K, f)$, define a bounded operator $S : H \to H$ by $S = diagonal(T_{v_1}, \ldots, T_{v_n})$. Then it is clear that $S(E_i) \subset E_i$ for $i = 1, \ldots, n$. Since $T_{r(\alpha_j)} f_{\alpha_j} = f_{\alpha_j} T_{s(\alpha_j)}$, $S(E_{n+j}) \subset E_{n+j}$ for $j = 1, \ldots m$. Hence $\varphi(T) = S$ define a homomorphism $\varphi : End(K, f) \to End\ \mathcal{S}$. Conversely Let S be in $End\ \mathcal{S}$. Since $S(E_i) \subset E_i$ for $i = 1, \ldots, n$, we can define bounded operators $T_{v_i} : K_{v_i} \to K_{v_i}$ such that $S = diagonal(T_{v_1}, \ldots, T_{v_n})$. Since $S(E_{n+j}) \subset E_{n+j}$ for $j = 1, \ldots m$, we have $T_{r(\alpha_j)} f_{\alpha_j} = f_{\alpha_j} T_{s(\alpha_j)}$. This shows that $T := (T_{v_k})_k \in End(K, f)$. Hence $\psi(S) = T$ define a homomorphism $\psi : End\ \mathcal{S} \to End(K, f)$. Since $\psi \circ \varphi = id$ and $\varphi \circ \psi = id$, $End(K, f) \cong End(\mathcal{S})$. □

Theorem 5.3. *Let $\Gamma = (V, E, s, r)$ be a finite quiver. Let (K, f) be a Hilbert representation of Γ. Then there exists a natural number n and a system $\mathcal{S}$ of n-subspaces such that $End(K, f) \cong End(\mathcal{S})$.*

Proof. Let $\Gamma = (V, E, s, r)$ be a finite quiver and (K, f) a Hilbert representation of Γ. Then there exists another finite quiver $\Gamma' = (V', E', s, r)$ without self-loops and a Hilbert representation (K', f') of Γ' such that $End(K, f) \cong End(K', f')$. In fact, replace "locally" each n-loop $\alpha_1, \ldots, \alpha_n : v \to v$ by $(n+1)$- Kronecker-like edges $\beta_1, \ldots, \beta_{n+1} : v \to v'$ to get Γ'. Any edge $\gamma : v \to w (\neq v)$ is also replaced by $\gamma' : v' \to w$. Then

Define (K', f') by $K'_v = K_v$ and $K'_{v'} = K_v$, and $f'_{\beta_1} = f_{\alpha_1}, \ldots, f'_{\beta_n} = f_{\alpha_n}$ and $f'_{\beta_{n+1}} = id$. We also put $f'_{\gamma'} = f_\gamma$. Then we have $End(K, f) \cong End(K', f')$. Next apply the lemma 5.2 for the finite quiver $\Gamma' = (V', E', s, r)$ without self-loops and a Hilbert representation (K', f') of Γ'. □

References

[Au] M. Auslander, *Large modules over artin algebras,* In: Algebra, Topology and Category Theory, Academic Press New York (1976), 1-17.

[BGP] I. N. Bernstein, I. M. Gelfand and V. A. Ponomarev, Coxeter functors and Gabriel's theorem, Russian Math. Surveys, 28 (1973), 17-32.

[DZ] A. Dean and F. Zorzitto, *A criterion for pure simplicity,* J.Algebra **132**, (1990), 50-71.

[DR] V. Dlab and C. M. Ringel, Indecomposable representations of graphs and algebras, Memoirs Amer. Math. Soc. 6 (1976), no. 173.

[DF] P. Donovan and M. R. Freislish, *The representation theory of finite graphs and associated algebras,* Carleton Math. Lect. Notes, Vol. 5, 1973, pp. 1-119.

[EW1] M. Enomoto and Y. Watatani, *Relative position of four subspaces in a Hilbert space,* Adv. Math. 201 (2006), 263-317.

[EW2] M. Enomoto and Y. Watatani, *Exotic indecomposable systems of four subspaces in a Hilbert space,* Integral Equations Operator Theory 59 (2007), 149-164.

[EW3] M. Enomoto and Y. Watatani, *Indecomposable representations of quivers on infinite-dimensional Hilbert spaces,* J. Funct. Anal. **256** (2009), 959-991.

[Fi] U. Fixman, *On algebraic equivalence between pairs of linear transformations,* Trans.Amer. Math. Soc. **113** (1964), 424-453.

[Ga] P. Gabriel, *Unzerlegbare Darstellungen I,* Manuscripta Math. 6 (1972), 71-103.

[GR] P. Gabriel and A.V. Roiter, Representations of Finite-Dimensional Algebras, Springer-Verlag, 1997.

[GP] I. M. Gelfand and V. A. Ponomarev, *Problems of linear algebra and classification of quadruples of subspaces in a finite-dimensional vector space,* Coll. Math. Spc. Bolyai 5, Tihany (1970), 163-237.

[Gi] F. Gilfeather, *Strong reducibility of operators,* Indiana Univ. Math. J. 22, (1972), 393-397.

[GHJ] F. Goodman, P. de la Harpe and V. Jones , Coxeter Graphs and Towers of Algebras, MSRI Publications, 14, Springer, Berlin, 1989.

[HRR] K.J. Harrison, H. Radjavi and P. Rosenthal, *A transitive medial subspace lattice*, Proc. Amer. Math. Soc. 28 (1971), 119-121.

[JW1] C.Jiang and Z. Wang, *Strongly Irreducible Operators on Hilbert Space*, Longman, 1998.

[JW2] C.Jiang and Z. Wang, *Structure of Hilbert Space Operators*, World Scientific, 2006.

[Ka] V. G. Kac, *Infinite root systems, representations of graphs and invariant theory,* Invent. Math. 56 (1980), 57-92.

[KR] H. Krause and C. M. Ringel, ed., *Infinite Length Modules,* Birkhäuser, 2000.

[KRo] S. A. Kruglyak and A. V. Roiter, *Locally scalar representations of graphs in the category of Hilbert spaces,* Funct. Anal. Appl. 39 (2005), 91-105.

[KRS] S. Kruglyak, V. Rabanovich and Y. Samoilenko, *On sums of projections,* Functional Anal. Appl., 36(2002), 182-195.

[MS] Y. P. Moskaleva and Y. S. Samoilenko, *Systems of* n *subspaces and representations of *-algebras generated by projections,* Methods Funct. Anal. Topology 12 (2006), no. 1, 57-73.

[Na1] L. A. Nazarova,*Representations of quadruples,* Izv. Akad. Nauk SSSR Ser. Mat. 31 (1967), 1361-1377.

[Na2] L. A. Nazarova, *Representation of quivers of infinite type,* Izv. Akad. Nauk SSSR, Ser. Mat., 37 (1973), 752-791.

[Ok] F. Okoh, *Applications of linear functional to Kronecker modules I,* Linear Algebra Appl.76 (1986), 165-204.

[RR] H. Radjavi and P. Rosenthal, *Invariant Subspaces*, Springer-Verlag, 1973.

[Ri1] C.M. Ringel, *Infinite dimensional representations of finite dimensional hereditary algebras*, Symposia Mathematica 23 (1979), Istituto Naz. Alta Matematica. 321-412.

[Ri2] C.M. Ringel, *Infinite length modules, Some examples as introduction,* Infinite Length Modules, Birkhäuser, Basel, 2000, 1-73.

[Sh] A. L. Shields, *Weighted shift operators and analytic function theory,* Topics in OperatorTheory, Math. Surveys Monographs, Amer. Math. Soc., Providence, RI 13 (1974), 49-128

(Masatoshi Enomoto) INSTITUTE OF EDUCATION AND RESEARCH, KOSHIEN UNIVERSITY, TAKARAZUKA, HYOGO 665-0006, JAPAN

(Yasuo Watatani) DEPARTMENT OF MATHEMATICAL SCIENCES, KYUSHU UNIVERSITY, MOTOOKA, FUKUOKA, 819-0395, JAPAN